

Indinavir (IDV, Crixivan)

For additional information see Drugs@FDA:

<http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm>

Formulations

Capsules: 100 mg, 200 mg, and 400 mg

Dosing Recommendations

Neonate/infant dose:

IDV is not approved for use in neonates/infants.

IDV should not be administered to neonates because of the risks associated with hyperbilirubinemia (kernicterus).

Pediatric dose:

IDV is not approved for use in children.

A range of IDV doses (234–500 mg/m² of body surface area) boosted by low-dose ritonavir (RTV) has been studied in children (see [Pediatric Use](#)).

Adolescent/adult dose:

800 mg IDV + 100 or 200 mg RTV every 12 hours.

Selected Adverse Events

- Nephrolithiasis
- Gastrointestinal (GI) intolerance, nausea
- **Hepatitis**
- Indirect hyperbilirubinemia
- Hyperlipidemia
- Headache, asthenia, blurred vision, dizziness, rash, metallic taste, thrombocytopenia, alopecia, and hemolytic anemia
- Hyperglycemia
- Fat maldistribution
- Possible increased bleeding episodes in patients with hemophilia

Special Instructions

- Administer IDV on an empty stomach 1 hour before or 2 hours after a meal (or administer with a light meal). When given in combination with RTV, meal restrictions are no longer necessary.
- Adequate hydration is required to minimize risk of nephrolithiasis (≥48 oz of fluid daily in adult patients).
- If coadministered with didanosine (ddI), give IDV and ddI ≥1 hour apart on an empty stomach.
- IDV capsules are sensitive to moisture; store at room temperature (59–86°F) in original container with desiccant.

Metabolism

- Cytochrome P450 3A4 (CYP3A4) inhibitor and substrate.
- **Dosing in patients with hepatic impairment:** Decreased dosage should be used in patients with mild-to-moderate hepatic impairment (recommended dose for adults is 600 mg IDV every 8 hours). No dosing information is

available for children with any degree of hepatic impairment or for adults with severe hepatic impairment.

Drug Interactions (See also the [Guidelines for the Use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents](#)):

- *Metabolism*: CYP3A4 is the major enzyme responsible for indinavir metabolism. There is potential for multiple drug interactions.
- Before indinavir is administered, the patient's medication profile should be carefully reviewed for potential drug interactions with indinavir.

Major Toxicities:

- *More common*: Nausea, abdominal pain, headache, metallic taste, dizziness, asymptomatic hyperbilirubinemia (10%), lipid abnormalities, pruritis, and rash. Nephrolithiasis/urolithiasis with indinavir crystal deposits.
- *Less common (more severe)*: Fat redistribution.
- *Rare*: New onset diabetes mellitus, hyperglycemia, ketoacidosis, exacerbation of pre-existing diabetes mellitus, spontaneous bleeding in hemophiliacs, acute hemolytic anemia, and hepatitis (life threatening in rare cases).
- *Pediatric specific*: The cumulative frequency of nephrolithiasis is higher in children (29%) than in adults (12.4%).

Resistance: The International Antiviral Society-USA (IAS-USA) maintains a list of updated resistance mutations (see http://www.iasusa.org/resistance_mutations/index.html) and the Stanford University HIV Drug Resistance Database offers a discussion of each mutation (see <http://hivdb.stanford.edu/pages/GRIP/IDV.html>).

Pediatric Use: Indinavir has not been approved by the Food and Drug Administration (FDA) for use in the pediatric population. Even though indinavir was one of the first protease inhibitors (PIs) to be studied in children, its use in pediatrics has never been common and is currently very rare¹.

Both unboosted and ritonavir-boosted indinavir have been studied in HIV-infected children. Data in children indicate that a dose of 500–600 mg of unboosted indinavir per meter² of body surface area given every 8 hours results in peak blood concentrations and areas under the curve (AUC) slightly higher than those in adults but considerably lower trough concentrations. A significant proportion of children have trough indinavir concentrations less than the 0.1 mg/L value associated with virologic efficacy in adults²⁻¹⁷. Studies in small groups of children of a range of ritonavir-boosted indinavir doses have shown that 500 mg indinavir per meter² of body surface area plus 100 mg ritonavir per meter² of body surface area twice daily is probably too high¹⁴, that 234–250 mg indinavir per meter² of body surface area plus low-dose ritonavir twice daily is too low¹⁸⁻¹⁹, and that 400 mg indinavir per meter² of body surface area plus 100–125 mg ritonavir per meter² of body surface area twice daily results in exposures approximating those seen with 800 mg indinavir/100 mg ritonavir twice daily in adults, albeit with considerable interindividual variability and high rates of toxicity^{2,5,19}.

As noted above, the cumulative frequency of nephrolithiasis is substantially higher in children (29%) than in adults (12.4%, range across clinical trials 4.7%–34.4%)²⁰. This is likely due to the difficulty in maintaining hydration adequate to minimize risk of nephrolithiasis in children. Finally, a large analysis of more than 2,000 HIV-infected children from PACTG 219 demonstrated a hazard ratio of 1.7 for the risk of renal dysfunction among children receiving combination antiretroviral therapy (cART) with indinavir²¹.

References

1. Van Dyke RB, Patel K, Siberry GK, et al. Antiretroviral Treatment of U.S. Children with Perinatally-Acquired HIV Infection: Temporal Changes in Therapy between 1991 and 2009 and Predictors of Immunologic and Virologic Outcomes. *J Acquir Immune Defic Syndr*. 2011;57(2):165-173.
2. Bergshoeff AS, Fraaij PL, van Rossum AM, et al. Pharmacokinetics of indinavir combined with low-dose ritonavir in human immunodeficiency virus type 1-infected children. *Antimicrob Agents Chemother*. 2004;48(5):1904-1907.
3. Burger DM, van Rossum AM, Huguenot PW, et al. Pharmacokinetics of the protease inhibitor indinavir in human immunodeficiency virus type 1-infected children. *Antimicrob Agents Chemother*. 2001;45(3):701-705.
4. Fletcher CV, Brundage RC, Remmel RP, et al. Pharmacologic characteristics of indinavir, didanosine, and stavudine in human immunodeficiency virus-infected children receiving combination therapy. *Antimicrob Agents Chemother*. 2000;44(4):1029-1034.
5. Fraaij PL, Bergshoeff AS, van Rossum AM, et al. Changes in indinavir exposure over time: a case study in six HIV-1-infected children. *J Antimicrob Chemother*. 2003;52(4):727-730.
6. Fraaij PL, Verweel G, van Rossum AM, et al. Indinavir/low-dose ritonavir containing HAART in HIV-1 infected children has potent antiretroviral activity, but is associated with side effects and frequent discontinuation of treatment. *Infection*. 2007;35(3):186-189.
7. Gatti G, Vigano A, Sala N, et al. Indinavir pharmacokinetics and pharmacodynamics in children with human immunodeficiency virus infection. *Antimicrob Agents Chemother*. 2000;44(3):752-755.
8. Jankelevich S, Mueller BU, Mackall CL, et al. Long-term virologic and immunologic responses in human immunodeficiency virus type 1-infected children treated with indinavir, zidovudine, and lamivudine. *J Infect Dis*. 2001;183(7):1116-1120.
9. Kline MW, Fletcher CV, Harris AT, et al. A pilot study of combination therapy with indinavir, stavudine (d4T), and didanosine (ddI) in children infected with the human immunodeficiency virus. *J Pediatr*. 1998;132(3 Pt 1):543-546.
10. Mueller BU, Sleasman J, Nelson RP, et al. A phase I/II study of the protease inhibitor indinavir in children with HIV infection. *Pediatrics*. 1998;102(1 Pt 1):101-109.
11. Pelton SI, Stanley K, Yogev R, et al. Switch from ritonavir to indinavir in combination therapy for HIV-1-infected children. *Clin Infect Dis*. 2005;40(8):1181-1187.
12. van Rossum AM, de Groot R, Hartwig NG, et al. Pharmacokinetics of indinavir and low-dose ritonavir in children with HIV-1 infection. *AIDS*. 2000;14(14):2209-2210.
13. van Rossum AM, Niesters HG, Geelen SP, et al. Clinical and virologic response to combination treatment with indinavir, zidovudine, and lamivudine in children with human immunodeficiency virus-1 infection: a multicenter study in the Netherlands. On behalf of the Dutch Study Group for Children with HIV-1 infections. *J Pediatr*. 2000;136(6):780-788.

14. van Rossum AM, Dieleman JP, Fraaij PL, et al. Persistent sterile leukocyturia is associated with impaired renal function in human immunodeficiency virus type 1-infected children treated with indinavir. *Pediatrics*. 2002;110(2 Pt 1):e19.
15. Verweel G, van Rossum AM, Hartwig NG, et al. Treatment with highly active antiretroviral therapy in human immunodeficiency virus type 1-infected children is associated with a sustained effect on growth. *Pediatrics*. 2002;109(2):E25.
16. Vigano A, Dally L, Bricalli D, et al. Clinical and immuno-virologic characterization of the efficacy of stavudine, lamivudine, and indinavir in human immunodeficiency virus infection. *J Pediatr*. 1999;135(6):675-682.
17. Wintergerst U, Hoffmann F, Solder B, et al. Comparison of two antiretroviral triple combinations including the protease inhibitor indinavir in children infected with human immunodeficiency virus. *Pediatr Infect Dis J*. 1998;17(6):495-499.
18. Plipat N, Cressey TR, Vanprapar N, et al. Efficacy and plasma concentrations of indinavir when boosted with ritonavir in human immunodeficiency virus-infected Thai children. *Pediatr Infect Dis J*. 2007;26(1):86-88.
19. Curras V, Hocht C, Mangano A, et al. Pharmacokinetic study of the variability of indinavir drug levels when boosted with ritonavir in HIV-infected children. *Pharmacology*. 2009;83(1):59-66.
20. FDA. Crixivan Label.
http://www.accessdata.fda.gov/drugsatfda_docs/label/2010/020685s073lbl.pdf. 2010.
21. Andiman WA, Chernoff MC, Mitchell C, et al. Incidence of persistent renal dysfunction in human immunodeficiency virus-infected children: associations with the use of antiretrovirals, and other nephrotoxic medications and risk factors. *Pediatr Infect Dis J*. 2009;28(7):619-625.